

A New Species of Eriophyid Mite Injurious to Tea Plant in Japan (Acari: Eriophyidae)

Fujio KADONO¹⁾

上遠野富士夫¹⁾: 日本の茶樹を加害するフシダニの1新種

Abstract A Japanese species of tea rust mite so far regarded as *Acaphylla theae* (WATT) or *A. steinwedeni* KEIFER is described under the name *A. theavagrans* sp. nov.

Since the anonymous report published in 1917, the Japanese tea rust mite was regarded as the same as *Acaphylla theae* (WATT, 1903) (= *Phytoptus theae* W.) originally described from India (e.g. MINAMIKAWA, 1950, 1957; MINAMIKAWA and OSAKABE, 1979; EHARA and KADONO, 1987). Recently, HUANG (1980) identified the same Japanese mite with *Acaphylla steinwedeni* KEIFER, 1943, known from California. However, I have come to the conclusion that the mite is neither *A. theae* nor *A. steinwedeni*, but a new species. It will be described and illustrated in this paper.

The abbreviations used here are the same as those explained in KADONO (1981).

Acaphylla theavagrans sp. nov.

[Japanese name: Cha-no-naga-sabidani]

(Fig. 1)

Phyllocoptes (*Phytoptus*) *theae* (nec WATT): anonymous, 1917, p. 122.

Phytoptus theae (nec WATT): MINAMIKAWA, 1950, p. 50.

Eriophyes theae (nec WATT): MINAMIKAWA, 1957, p. 153.

Acaphylla theae (nec WATT): MINAMIKAWA & OSAKABE, 1979, p. 236, fig. 101, p. 280; EHARA & KADONO, 1987, p. 23.

Acaphylla steinwedeni (nec KEIFER): HUANG, 1980, p. 291, pl. 133-C.

Female. Body fusiform, orange to orange yellow in color. Rostrum curved ventrally; antepical seta forked apically, 17 μ m long. Shield subtriangular in anterior outline; anterior lobe over rostral base; median line absent; admedian lines running in zigzag from anterior shield lobe to a point slightly posterior to

1) Chiba Prefectural Agricultural Experiment Station, 808, Daizenno-cho, Midori-ku, Chiba, 266 Japan

千葉県農業試験場

Accepted November 10, 1992

the transverse level through shield setae, and curving inwardly each other at rear; transverse lines running from admedian lines at anterior 1/3 to lateral shield margin; irregularly curved lines present in the area surrounded by admedian lines and transverse lines from admedian lines; dorsal tubercles present ahead of rear shield margin; dorsal seta minute, directing upwardly. Foreleg without 1st coxal setae; tibial seta 4 μm long, arising from proximal 1/4 of the segment; claw strongly curved, with a large knob terminally; featherclaw divided deeply, with 3-rayed. Hindleg without patellar setae. Coxae almost smooth; first setiferous coxal tubercles absent; second coxal tubercles located at level before third tubercles; coxae touching each other. Abdomen with 30–34 tergites and 69–79 sternites; tergites smooth; sternites with microtubercles.

Genitalia 18 μm long, 25 μm wide; coverflap with 6–10 ribs anteriorly, with microtubercles basally. Seta *ls* on sternite 11–14, *vs*₁ on sternite 28–33, *vs*₂ on sternite 46–52, *vs*₃ on sternite 6 from rear. Mean (range in parentheses) in μm , length unless otherwise stated ($n=10$): body 209 (201–215), width 77 (68–85); shield 59, width 77; rostrum 31; foreleg 37, femur 11, patella 6, tibia 8, tarsus 8, claw 6, featherclaw 5; hindleg 35, femur 9, patella 5, tibia 7, tarsus 7, claw 5, featherclaw 5; setae *ds* 4, *ls* 14, *vs*₁ 53, *vs*₂ 43, *vs*₃ 20, *cxs*₂ 12, *cxs*₃ 31, *gs* 12, *acs* missing; intervals between setae: *ds*–*ds* 17, *ls*–*ls* 54, *vs*₁–*vs*₁ 28, *vs*₂–*vs*₂ 14, *vs*₃–*vs*₃ 25, *cxs*₂–*cxs*₂ 9, *cxs*₃–*cxs*₃ 25, *gs*–*gs* 15.

Male. Similar to female except for genitalia. Abdomen with 29–30 tergites and 58–65 sternites; seta *ls* on sternite 9–12, *vs*₁ on sternite 21–25, *vs*₂ on sternite 33–41, *vs*₃ on sternite 6 from rear. Mean (range in parentheses) in μm , length unless otherwise stated ($n=2$): body 163 (162–164), width 64 (63–66); shield 47, width 64; genitalia 17, width 16.

Type series. Holotype: ♀, paratypes: 9♀♀ and 2♂♂, Kasai, Hyogo Pref., 10–VI–1992 (K. KASAMATSU), on the tea plant, *Thea sinensis* L. The type series is deposited in the collection of Chiba Prefectural Agricultural Experiment Station.

Distribution. Japan (Honshu); Taiwan.

Remarks. *Acaphylla theavagrans* sp. nov. is similar to *A. steinwedeni* KEIFER, 1943 (California), but differs from the latter in the shield having the apparent transverse lines from the admedian lines to the lateral shield margin and some curved lines outside the admedians. This new species also has a resemblance to *A. theae* (WATT, 1903) (WATT & MANN, 1903; India). However, judging from the original description of *A. theae*, the new mite differs from *theae* in having no rounded elevations on the central dorsum of the abdomen. An *Acaphylla* species living on tea plant in Taiwan was referred to *A. steinwedeni* by HUANG (1974). However, the Taiwanese mite should be assigned to *A. theavagrans* sp. nov.

Acknowledgments

I wish to express my heartfelt thanks to Professor Dr. S. EHARA, Tottori University, for his helpful comments on the draft of the paper. I also thank Dr. T. HAMAMURA, National Research Institute of Vegetable, Ornamental Plants and Tea, and Mr. S. KUBOTA, Shizuoka Tea Experiment Station for information about the new mite.




Fig. 1. *Acaphylla theavagrans* sp. nov.—A, Body, dorsal view, ds, dorsal setae; B, body, lateral view; C, side skin of abdomen; D, coxae and genitalia; E, internal genital structure; F, 1st leg; Fa, 2nd leg; G, featherclaw.

摘 要

これまで *Acaphylla theae* (WATT) あるいは *A. steinwedeni* KEIFER と同定されてきたチャノナガサビダニが新種 *A. theavagrans* KADONO として記載された。

References

- ANONYM, 1917. Annual Report of Shizuoka Agricultural Experiment Station. 134 pp. (In Japanese.)
 EHARA, S., & F. KADONO, 1987. Acarina. In Jpn. Soc. Appl. Entomol. Zool. (ed.): *Major Insect and Other Pests of Economic Plants in Japan*, pp. 21–25. Jpn. Plant Protec.

- Assoc., Tokyo. (In Japanese.)
- HUANG, T., 1974. Records of six eriophyid mites associated with economic plants in Taiwan. *J. Agr. For.*, **23**: 75-88.
- 1980. *Acaphylla steinwedeni* KEIFER. In EHARA, S. (ed.): *Illustrations of the Mites and Ticks of Japan*, p.291. Zenkoku Noson Kyoiku Kyokai, Tokyo. (In Japanese.)
- KADONO, F., 1981. Two new eriophyid mites from Japan (Acarina: Eriophyidae). *Appl. Entomol. Zool.*, **16**: 419-422.
- KEIFER, H. H., 1943. Eriophyid mites XIII. *Bull. Calif. Dept. Agr.*, **32**: 212-222.
- MINAMIKAWA, J., 1950. A list of the tea injurious insects in Japan. *Chagyo Gijutsu Kenkyu*, (3): 47-50. (In Japanese.)
- 1957. A list of the tea injurious insects in Japan. *Botyu-Kagaku*, **22**: 149-154. (In Japanese.)
- & M.OSAKABE, 1979. Insect Pests of Tea Plant in Japan. iv+322 pp. Jpn. Plant Protec. Assoc., Tokyo. (In Japanese.)
- WATT, G., & H. H. MANN, 1903. The Pests and Blights of the Tea Plant (2nd ed.). 416 pp. Office of Supt. Govt. Printing, Calcutta.